

Bijlage: Literatuurverwijzingen

PEMF en TMS

Psyche

Blumberger DM, Mulsant BH, Fitzgerald PB, et al. A randomized double-blind sham-controlled comparison of unilateral and bilateral repetitive transcranial magnetic stimulation for treatment-resistant major depression. *World J Biol Psychiatry*. July 2011.

Lappin MS, Lawrie FW, Richards TL, Kramer ED. Effects of pulsed electromagnetic therapy on multiple sclerosis fatigue and quality of life: a double-blind, placebo controlled trial. *Altern Ther Health Med*. July-August 2003;9(4):38-48.

Demitrack MA, Thase ME. Clinical significance of transcranial magnetic stimulation (TMS) in the treatment of pharmacoresistant depression: synthesis of recent data. *Psychopharmacol Bull*. 2009;42(2):5-38.

Poulet E, Haesebaert F, Saoud M, Suaud-Chagny MF, Brunelin J. Treatment of schizophrenic patients and rTMS. *Psychiatr Danub*. November 2010;22 Suppl 1:S143-6.

Watts BV, Landon B, Groft A, Young-Xu Y. A sham controlled study of repetitive transcranial magnetic stimulation for posttraumatic stress disorder. *Brain Stimul*. January 2012;5(1):38-43.

Blom RM, Figee M, Vulink N, Denys D. Update on repetitive transcranial magnetic stimulation in obsessive-compulsive disorder: different targets. *Curr Psychiatry Rep*. August 2011;13(4):289-94.

Bentwich J, Dobronevsky E, Aichenbaum S. Beneficial effect of repetitive transcranial magnetic stimulation combined with cognitive training for the treatment of Alzheimer's disease: a proof of concept study. *J Neural Transm*. March 2001;118(3):463-71.

Pal E, Nagy F, Aschermann Z, Balazs, E, Kovacs N. The impact of left prefrontal repetitive transcranial magnetic stimulation on depression in Parkinson's disease: a randomized, double-blind, placebo-controlled study. *Mov Disord*. October 2010;25(14):2311-7.

Martiny. K, Lunde. M, Bech. P, (2010). Transcranial Low Voltage Pulsed Electromagnetic Fields in Patients with Treatment-Resistant Depression. *Biol Psychiatry* 2010;68:163–169

Pascual-Leone A, Rubio B, Pallardo F, Catala MD. Rapid-rate transcranial magnetic stimulation of the left dorsolateral prefrontal cortex in drug-resistant depression. Lancet 1996; 347: 233-237.

Tergau F, Naumann U, Paulus W, Steinhoff BJ. Low-frequency repetitive transcranial magnetic stimulation improves intractable epilepsy. Lancet 1999; 353: 2209.

Hoffman RE, Boutros NN, Hu S, Berman RM, Krystal JH, Charney DS. Transcranial magnetic stimulation and auditory hallucinations in schizophrenia. Lancet 2000; 355: 1073-1075.

Kortekaas R, van Nierop LE, Baas VG, Konopka K-H, Harbers M, et al. (2013) A Novel Magnetic Stimulator Increases Experimental Pain Tolerance in Healthy Volunteers - A Double-Blind Sham-Controlled Crossover Study. PLoS ONE 8(4): e61926. doi:10.1371/journal.pone.0061926

Bewegingsapparaat

Ciombor DM1, Aaron RK, Wang S, Simon B: Modification of osteoarthritis by pulsed electromagnetic field--a morphological study. Osteoarthritis Cartilage. 2003 Jun;11(6):455-62.

Siskin B F, Walker J. Therapeutic aspects of electromagnetic fields for soft-tissue healing. In Blank M ed. Electromagnetic fields: Biological interactions and Mechanisms. Advances in Chemistry Series. Vol 250. Washington, DC: American Chemical Society; 1995:277-285.

Lee PB, Kim YC, Lim YJ, et al. Efficacy of pulsed electromagnetic therapy for chronic lower back pain: a randomized, double-blind, placebocontrolled study. J Int Med Res. March-April 2006;34(2):160-7.

Thomas AW, Graham K, Prato FS, et al. A randomized, double-blind, placebo-controlled trial using low-frequency magnetic fields in the treatment of musculoskeletal chronic pain. Pain Res Manag. Winter 2007;12(4):249-58.

Sutbeyaz ST, Sezer N, Koseoglu BF. The effect of pulsed electromagnetic fields in the treatment of cervical osteoarthritis: a randomized, double-blind, sham-controlled trial. Rheumatol Int. February 2006;26(4): 320-4.

Nicolakis P, Kollmitzer J, Crevenna R, Bittner C, Erdoganmus CB, Nicolakis J. Pulsed magnetic field therapy for osteoarthritis of the knee – a double-blind, sham-controlled trial. Wien Klin Wochenschr. August 2002; 114(15-16):678-84.

Uzunca K, Birtane M, Tastekin N. Effectiveness of pulsed electromagnetic field therapy in lateral epicondylitis. *Clin Rheumatol*. January 2007;26(1):69-74.

Benazzo F, Zanon G, Pederzini L, et al. Effects of biophysical stimulation in patients undergoing arthroscopic reconstruction of anterior cruciate ligament: prospective, randomized and double blind study. *Knee Surg Sports Traumatol Arthrosc*. June 2008;16(6):595-601.

Mooney V. A randomized double-blind prospective study of the efficacy of pulsed electromagnetic fields for interbody lumbar fusions. *Spine* July 1990;15(7):708-12.

Binder A, Parr G, Hazleman B, Fitton-Jackson S. Pulsed electromagnetic field therapy of persistent rotator cuff tendinitis. A double-blind controlled assessment. *Lancet*. March 1984;1(8379):695-8.

Grant G, Cadossi R, Steinberg G. Protection against focal cerebral ischemia following exposure to pulsed electromagnetic field. *Bioelectromagnetics*. 1994;15(3):205-16.

Rubin CT, McLeod KJ, Lanyon LE. Prevention of osteoporosis by pulsed electromagnetic fields. *J Bone Joint Surg Am*. March 1989;71(3):411-7.

Tabrah F, Hoffmeier M, Gilbert F Jr, Batkin S, Bassett CA. Bone density changes in osteoporosis-prone women exposed to pulsed electromagnetic fields (PEMFs). *J Bone Miner Res*. May 1990;5(5):437-42.

Shen WW, Zhao JH. Pulsed electromagnetic fields stimulation affects BMD and local production of rats with disuse osteoporosis. *Bioelectromagnetics*. February 2010;31(2):113-19.

Jing D, Cai J, Shen G, et al. The preventive effects of pulsed electromagnetic fields on diabetic bone loss in streptozocin-treated rats. *Osteoporos Int*. June 2011;22(6):1885-95.

Celmetabolisme

Schnoke M, Midura RJ. Pulsed electromagnetic fields rapidly modulate intracellular signaling events in osteoblastic cells: comparison to parathyroid hormone and insulin. *J Orthop Res*. July 2007;25(7):933-40.

Mert T, Gunay I, Ocal I. Neurobiological effects of pulsed magnetic field on diabetes-induced neuropathy. *Bioelectromagnetics*. January 2010;31(1):39-47.

Luo E, Shen G, Xie K, et al. Alimentary hyperlipidemia of rabbits is affected by exposure to low-intensity pulsed magnetic fields. *Bioelectromagnetics*. December 2007;28(8):608-14.

Muehsam D, Lalezari P, Lekhraj R, Abruzzo P, Bolotta A, Marini M, Bersani F, Aicardi G, Pilla A, Casper Non-thermal radio frequency and static magnetic fields increase rate of hemoglobin deoxygenation in a cell-free preparation. *DPLoS One*. 2013 Apr 12;8(4):e61752. doi: 10.1371/journal.pone.0061752. Print 2013.

Immunité et résistance

Andrea Cossarizza, Stefano Angioni, Felice Petraglia, Andrea R. Genezzani, Daniela Monti, Miriam Capri, Ferdinando Bersani, Ruggero Cadossi, Claudio Franceschi: Exposure to Low Frequency Pulsed Electromagnetic Fields Increases Interleukin-1 and Interleukin-6 Production by Human Peripheral Blood Mononuclear Cells. *Experimental Cell Research* Volume 204, Issue 2, February 1993, Pages 385–387

Zafer Akan, Burak Aksu, Aysin Tulunay, Serpil Bilsel, Ayse Inhan-Garip: Extremely low-frequency electromagnetic fields affect the immune response of monocyte-derived macrophages to pathogens. *Bioelectromagnetics* Volume 31, Issue 8, pages 603–612, December 2010

Li JK, Lin JC, Liu HC, Chang WH. Cytokine release from osteoblasts in response to different intensities of pulsed electromagnetic field stimulation. *Electromagn Biol Med*. 2007;26(3):153-65.

Obermeier A, Matl FD, Friess W, Stemberger A Growth inhibition of *Staphylococcus aureus* induced by low-frequency electric and electromagnetic fields. *Bioelectromagnetics*. May 2009;30(4):270-9.

Matl FD, Obermeier A, Zlotnyk J, Friess W, Stemberger A, Burgkart R. Augmentation of antibiotic activity of low-frequency electric and electromagnetic fields examining *Staphylococcus aureus* in broth media. *Bioelectromagnetics*. July 2011;32(5):367-77.

Di Campli E, Di Bartolomeo S, Grande R, Di Giulio M, Cellini L. Effects of extremely low-frequency electromagnetic fields on *Helicobacter pylori* biofilm. *Curr Microbiol*. June 2010;60(6):412-8.

Pickering SA, Bayston R, Scammell BE. Electromagnetic augmentation of antibiotic efficacy in infection of orthopedic implants. *J Bone Joint Surg Br*. May 2003;85(4):588-93.

Zeng F, Zheng C, Zhang X, et al. Experimental studies on ultralow frequency pulsed gradient magnetic field inducing apoptosis of cancer cell and inhibiting growth of cancer cell. *Sci China C Life Sci*. February 2002 45(1):33-9.

Yamaguchi S, Ogiue-Ikeda M, Sekino M, Ueno S. Effects of pulsed magnetic stimulation on tumor development and immune function in mice. *Bioelectromagnetics*. January 2006;27(1):64-72.

Rihova B, Etrych T, Sirova M, Tomala J, Ulbrich K, Kovar M. Synergistic effect of EMF-BEMER-type pulsed weak electromagnetic field and HPMA-bound doxorubicin on mouse EL4 T-cell lymphoma. *J Drug Target*. December 2011;19(10):890-9.

Radeva M, Berg H. Differences in lethality between cancer cells and human lymphocytes caused by LF-electromagnetic fields. *Bioelectromagnetics*. October 2004;25(7):503-7.

Lichttherapie

Psyche

Meesters. Y, Waslander. M, (2009). Burnout and light treatment. *Stress & Health*, DOI:10.1002/smj.1250

Golden RN, Gaynes BN, Ekstrom RD, Hamer RM, Jacobsen FM, Suppes T, Wisner KL, Nemeroff CB. The efficacy of light therapy in the treatment of mood disorders: a review and meta-analysis of the evidence. *Am J Psychiatry*. 2005 Apr;162(4):656-62

Yasukouchi A, Hazama T, Kozaki T,. Variations in the light-induced suppression of nocturnal melatonin with special reference to variations in the pupillary light reflex in humans. *J Physiol Anthropol*. 2007 Mar;26(2):113-21.

Smolders K C H J, de Kort Y A W, Cluitmans P J M,. A higher illuminance induces alertness even during office hours: Findings on subjective measures, task performance and heart rate measures. *Physiology & Behavior* 107 (2012) 7–16

Rheaume YL, Manning BC, Harper DG, Volicer L. Effect of light therapy upon disturbed behaviors in Alzheimer patients. *American Journal of Alzheimer's Disease* 1998;13(6):291-295.

Van Someren EJ, Kessler A, Mirmiran M, Swaab DF. Indirect bright light improves circadian rest-activity rhythm disturbances in demented patients. *Biological Psychiatry* 1997;41(9): 955-963.

Riemersma-van der Lek RF, Swaab DF, Twisk J, Hol EM, Hoogendoorn WJG, van Someren EJW. Effect of bright light and melatonin on cognitive and noncognitive function in elderly residents of group care facilities. A randomized controlled trial. *The Journal of the American Medical Association* 2008;299(22): 2642-2655.

Sloane PD, Williams CS, Mitchell CM, Preisser JS, Wood W, Barrick AL, Hickman SE, Gill KS, Connell BR, Edinger J, Zimmerman S. High-intensity environmental light in dementia: effect on sleep and activity. *Journal of the American Geriatrics Society* 2007;55(10): 1524-1533

Van Hoof, J., Aarts, M.P.J., Rense, C.G., Schoutens, A.M.C.. Ambient bright light in dementia: Effects on behaviour and circadian rhythmicity *Building and Environment* 44, (2009) 146 - 155

Van Hoof, J., Schoutens, A.M.C., Aarts, M.P.J. High colour temperature lighting for institutionalised older people with dementia. *Building and Environment* 44(2009) 1959 – 1969

Domien Beersma, Luc Schlangen, Toine Schoutens. Invloed van licht op gezondheid onderschat. *Medisch Contact* Nr. 33/34 – pagina's 1658-1660, 15 augustus 2013