

Referenties:

- 1: Penn L, White M, Oldroyd J, Walker M, Alberti KG, Mathers JC. Prevention of type 2 diabetes in adults with impaired glucose tolerance: the European Diabetes Prevention RCT in Newcastle upon Tyne, UK. *BMC Public Health*. 2009 Sep 16;9:342.
- 2: Penn L, White M, Lindström J, den Boer AT, Blaak E, Eriksson JG, Feskens E, Ilanne-Parikka P, Keinänen-Kiukaanniemi SM, Walker M, Mathers JC, Uusitupa M, Tuomilehto J. Importance of weight loss maintenance and risk prediction in the prevention of type 2 diabetes: analysis of European Diabetes Prevention Study RCT. *PLoS One*. 2013;8(2):e57143.
- 3: den Boer AT, Herraets IJ, Stegen J, Roumen C, Corpeleijn E, Schaper NC, Feskens E, Blaak EE. Prevention of the metabolic syndrome in IGT subjects in a lifestyle intervention: Results from the SLIM study. *Nutr Metab Cardiovasc Dis*. 2013 Feb 22.
- 4: Gray LJ, Cooper N, Dunkley A, Warren FC, Ara R, Abrams K, Davies MJ, Khunti K, Sutton A. A systematic review and mixed treatment comparison of pharmacological interventions for the treatment of obesity. *Obes Rev*. 2012 Jun;13(6):483-98.
- 5: Lakerveld J, Bot SD, Chinapaw MJ, van Tulder MW, Kostense PJ, Dekker JM, Nijpels G. Motivational interviewing and problem solving treatment to reduce type 2 diabetes and cardiovascular disease risk in real life: a randomized controlled trial. *Int J Behav Nutr Phys Act*. 2013 Apr 19;10:47.
- 6: Dunkley AJ, Charles K, Gray LJ, Camosso-Stefinovic J, Davies MJ, Khunti K. Effectiveness of interventions for reducing diabetes and cardiovascular disease risk in people with metabolic syndrome: systematic review and mixed treatment comparison meta-analysis. *Diabetes Obes Metab*. 2012 Jul;14(7):616-25.
- 7: Eakin EG, Reeves MM, Winkler E, Healy GN, Dunstan DW, Owen N, Marshal AM, Wilkie KC. Six-Month Outcomes from Living Well with Diabetes: A Randomized Trial of a Telephone-Delivered Weight Loss and Physical Activity Intervention to Improve Glycemic Control. *Ann Behav Med*. 2013 Apr 23. [Epub ahead of print]
- 8: Murrock CJ, Taylor E, Marino D. Dietary challenges of managing type 2 diabetes in African-American women. *Women Health*. 2013;53(2):173-84.
- 9: Gilstrap LG, Malhotra R, Peltier-Saxe D, Slicas D, Pineda E, Culhane-Hermann C, Cook N, Fernandez-Golarz C, Wood M. Community-based primary prevention programs decrease the rate of metabolic syndrome among socioeconomically disadvantaged women. *J Womens Health (Larchmt)*. 2013 Apr;22(4):322-9. doi:

10.1089/jwh.2012.3854. Epub 2013 Mar 29.

10: Taylor R. Type 2 diabetes: etiology and reversibility. *Diabetes Care*. 2013 Apr;36(4):1047-55.

11: Taylor R. Banting Memorial lecture 2012: reversing the twin cycles of type 2 diabetes. *Diabet Med*. 2013 Mar;30(3):267-75.

12: Steven S, Lim EL, Taylor R. Population response to information on reversibility of Type 2 diabetes. *Diabet Med*. 2013 Apr;30(4):e135-8.

13: <http://www.ajmc.com/publications/issue/2013/2013-1-vol19-n3/effectiveness-and-cost-effectiveness-of-diabetes-prevention-among-adherent-participants/5>

14: The Look AHEAD Research Group: Cardiovascular Effects of Intensive Lifestyle Intervention in Type 2 Diabetes. *N Engl J Med* 2013; 369:145-154

15: Hans Wahrenberg Katarina Hertel, Britt-Marie Leijonhufvud, Lars-Göran Persson, Eva Toft, Peter Arner: Use of waist circumference to predict insulin resistance: retrospective study *BMJ* 2005;330:1363

16: Logtenberg SJ1, Kleefstra N, Ubink-Veltmaat LJ, Houweling ST, Bilo HJ. Intensification of therapy and no increase in body mass index with longer disease duration in type 2 diabetes mellitus (ZODIAC-5). *Fam Pract*. 2007 Dec;24(6):529-31. Epub 2007 Nov 26.

17: Booz&Co / Novo Nordisk. *Diabetes Care in The Netherlands: improving health and wealth*. 2011

18: Jazet IM1, Ouwens DM, Schaart G, Pijl H, Keizer H, Maassen JA, Meinders AE. Effect of a 2-day very low- energy diet on skeletal muscle insulin sensitivity in obese type 2 diabetic patients on insulin therapy. *Metabolism*. 2005 Dec;54(12):1669-78.

19: Jazet IM1, Pijl H, Frölich M, Romijn JA, Meinders AE. Two days of a very low calorie diet reduces endogenous glucose production in obese type 2 diabetic patients despite the withdrawal of blood glucose- lowering therapies including insulin. *Metabolism*. 2005 Jun;54(6):705-12.

20. *Farmacotherapeutisch kompas*. Beschikbaar via <http://www.farmacotherapeutischkompas.nl> (referentiewaarden klinisch chemie) Geraadpleegd 7 mei 2014

21. Langeveld M, de Vries JH. Het magere resultaat van diëten. *Ned Tijdsch Geneesk* 2013;157: A6017

22. NHG standaard Diabetes Mellitus type 2, 2013

23. Risico's van verhoogd cholesterol. Hartstichting. Beschikbaar via <https://www.hartstichting.nl/risicofactoren/hog-cholesterol>. Geraadpleegd 4 juli 2014

24. NHG standaard (CVRM) Cardio Vasculair Risico Management. 2012

25. Evidence- based Voedingsrichtlijn voor diabetes type 1 en 2. Nederlandse Diabetes Federatie (NDF), Amersfoort, augustus 2010.

26. Metformine, SU-derivaten en Insuline. Beschikbaar via http://www.apotheek.nl/Medische_informatie/Medicijnen/Producten/ Geraadpleegd op 4 augustus 2014